arXiv:math/0105229v1 [math.QA] 28 May 2001

ON DERIVATION DEVIATIONS IN AN ABSTRACT PRE-OPERAD

Liivi Kluge and Eugen Paal

Abstract. We consider basic algebraic constructions associated with an *abstract* pre-operad, such as a $\smile$-algebra, total composition $\bullet$, *pre-coboundary* operator δ and *tribraces* $\{\cdot,\cdot,\cdot\}$. A derivation deviation of the pre-coboundary operator over the tribraces is calculated in terms of the $\smile$-multiplication and total composition.

Classification (MSC2000). 18D50.

Key words. Comp(osition), (pre-)operad, Gerstenhaber theory, cup, pre-coboundary, (tri)braces, derivation deviation.

1. Introduction and outline of the paper

We consider basic algebraic constructions associated with an *abstract* pre-operad C (Sec. 2), such as a $\smile$-algebra (Sec. 3), total composition $\bullet$ (Sec. 4), *pre-coboundary* operator δ (Sec. 5) and *tribraces* $\{\cdot,\cdot,\cdot\}$ (Sec. 6). Main result of the present paper is the Main Theorem (Sec. 6). By defining (see degree notations in Sec. 2) a *derivation deviation* (Definition 6.2) of the pre-coboundary operator δ over the tribraces $\{\cdot,\cdot,\cdot\}$,

$$\operatorname{dev}_{\{\cdot,\cdot,\cdot\}}\delta\,(h\otimes f\otimes g) := \delta\{h,f,g\} - \{h,f,\delta g\} - (-1)^{|g|}\{h,\delta f,g\} - (-1)^{|g|+|f|}\{\delta h,f,g\},$$

the Main Theorem states that in a the pre-operad C one has

$$(-1)^{|g|}\operatorname{dev}_{\{\cdot,\cdot,\cdot\}}\delta\,(h\otimes f\otimes g) = (h\bullet f)\smile g + (-1)^{|h|f} f\smile(h\bullet g) - h\bullet(f\smile g).$$

This formula (Main Theorem) modifies and generalizes Gerstenhaber's Theorem 5 from [2]. One can see the appearance of the extra terms with $\delta h, \delta f$ and δg inside the tribraces. The Gerstenhaber formula is obtained when arguments in tribraces $\{h,f,g\}$ are the Hochshild *cocycles*, i. e. if $\delta h = \delta f = \delta g = 0$. As it was stated in [5, 14], on the Hochschild *cochain* level the extra terms with $\delta h, \delta f$ and δg also appear when modifying the Gerstenhaber formula. We work out these extra terms for an *abstract* pre-operad and give their interpretation via a *derivation deviation* of δ over the tribraces.

In this paper, we do not assume *associativity* constraint (see Example 5.2).

---

math.QA/0105229

Research supported in part by the ESF grant 3654.

2. Pre-operad (composition system)

Let K be a unital commutative associative ring, and let C^n ($n \in \mathbb{N}$) be unital K-modules. For *homogeneous* $f \in C^n$, we refer to n as the *degree* of f and often write (when it does not cause confusion) f instead of $\deg f$; for example, $(-1)^f := (-1)^n$, $C^f := C^n$ and $\circ_f := \circ_n$. Also, it is convenient to use the shifted (*desuspended*) degree $|f| := n-1$. Throughout this paper, we assume that $\otimes := \otimes_K$.

Definition 2.1. A linear (right) *pre-operad* (*composition system*) with coefficients in K is a sequence $C := \{C^n\}_{n\in\mathbb{N}}$ of unital K-modules (an $\mathbb{N}$-graded K-module), such that the following conditions hold:

1. For $0 \leq i \leq m-1$ there exist *partial compositions*
$$\circ_i \in \operatorname{Hom}(C^m \otimes C^n, C^{m+n-1}), \qquad |\circ_i| = 0.$$
2. For all $h \otimes f \otimes g \in C^h \otimes C^f \otimes C^g$, the *composition relations* hold,
$$(h \circ_i f) \circ_j g = \begin{cases} (-1)^{|f||g|}(h \circ_j g) \circ_{i+|g|} f & \text{if } 0 \leq j \leq i-1, \\ h \circ_i (f \circ_{j-i} g) & \text{if } i \leq j \leq i+|f|, \\ (-1)^{|f||g|}(h \circ_{j-|f|} g) \circ_i f & \text{if } i+f \leq j \leq |h|+|f|. \end{cases}$$
3. There exists a unit $\mathrm{I} \in C^1$ such that
$$\mathrm{I} \circ_0 f = f = f \circ_i \mathrm{I}, \qquad 0 \leq i \leq |f|.$$

In the 2nd item, the *first* and *third* parts of the defining relations turn out to be equivalent.

Remarks 2.2. A pre-operad is also called a *comp(osition) algebra* or *asymmetric operad* or *non-symmetric operad* or *non-Σ operad*. The concept of (*symmetric*) *operad* was formalized by May [9] as a tool for the theory of iterated loop spaces. Recent studies and applications can be found in [7].

Above we modified the Gerstenhaber *comp algebra* defining relations [1, 4] by introducing the sign $(-1)^{|f||g|}$ in the defining relations of the pre-operad. The modification enables us to keep track of (control) sign changes more effectively. One should also note that (up to sign) our $\circ_i$ is Gerstenhaber's $\circ_{i+1}$ from [4, 1]; we use the original (non-shifted) convention from [2, 3].

Example 2.3 (endomorphism pre-operad [2, 3, 4])**.** Let A be a unital K-module and $\mathcal{E}^n_A := \mathcal{E}nd^n_A := \operatorname{Hom}(A^{\otimes n}, A)$. Define the partial compositions for $f \otimes g \in \mathcal{E}^f_A \otimes \mathcal{E}^g_A$ as
$$f \circ_i g := (-1)^{i|g|} f \circ (\mathrm{id}_A^{\otimes i} \otimes g \otimes \mathrm{id}_A^{\otimes(|f|-i)}), \qquad 0 \leq i \leq |f|.$$
Then $\mathcal{E}_A := \{\mathcal{E}^n_A\}_{n\in\mathbb{N}}$ is a pre-operad (with the unit $\mathrm{id}_A \in \mathcal{E}^1_A$) called the *endomorphism pre-operad* of A. A few examples (without the sign factor) can be found in [3, 4] as well. We use the original indexing of [2, 3] for the defining formulae.

Example 2.4 (associahedra)**.** A geometrical example of a pre-operad (without signs $(-1)^{\cdots}$ in the defining relations) is provided by the Stasheff *associahedra*, which was first constructed in [11]. Quite a surprising realization of the associahedra as *truncated simplices* was discovered and studied in [10, 12, 8].

Notations 2.5 (scope of a pre-operad)**.** The *scope* of $(h \circ_i f) \circ_j g$ is given by

$$0 \le i \le |h|, \qquad 0 \le j \le |f| + |h|.$$

It follows from the defining relations of a pre-operad that the scope is a disjoint union of

$$\begin{aligned} B &:= \{(i,j) \in \mathbb{N} \times \mathbb{N} \,|\, 1 \le i \le |h| \,;\, 0 \le j \le i-1\}, \\ A &:= \{(i,j) \in \mathbb{N} \times \mathbb{N} \,|\, 0 \le i \le |h| \,;\, i \le j \le i+|f|\}, \\ G &:= \{(i,j) \in \mathbb{N} \times \mathbb{N} \,|\, 0 \le i \le |h|-1 \,;\, i+f \le j \le |f|+|h|\}. \end{aligned}$$

Note that the triangles B and G are symmetrically situated with respect to the parallelogram A in the scope $BAG := B \sqcup A \sqcup G$. The (recommended and impressive) picture is left for a reader as an exercise.

Recapitulation 2.6. The defining relations of a pre-operad can be easily rewritten as follows:

$$(h \circ_i f) \circ_j g = \begin{cases} (-1)^{|f||g|} (h \circ_j g) \circ_{i+|g|} f & \text{if } (i,j) \in B, \\ h \circ_i (f \circ_{j-i} g) & \text{if } (i,j) \in A, \\ (-1)^{|f||g|} (h \circ_{j-|f|} g) \circ_i f & \text{if } (i,j) \in G. \end{cases}$$

The *first* (B) and *third* (G) parts of the relations turn out to be equivalent.

3. Cup

In this section, we recollect from [6] basic facts about cup-multiplication in an *abstract* pre-operad.

Definition 3.1 (cf. [4, 1])**.** In a pre-operad C, let $\mu \in C^2$. Define $\smile := \smile_\mu \colon C^f \otimes C^g \to C^{f+g}$ by

$$f \smile g := (-1)^f (\mu \circ_0 f) \circ_f g \in C^{f+g}, \qquad |\smile| = 1, \qquad f \otimes g \in C^f \otimes C^g.$$

Note that $\circ_f := \circ_{\deg f}$ and $|\smile| = 1$ means that $\deg \smile = 0$. The pair $\operatorname{Cup} C := \{C, \smile\}$ is called a $\smile$-algebra of C.

Example 3.2. For the endomorphism pre-operad (Example 2.3) $\mathcal{E}_A$, one has

$$f \smile g = (-1)^{fg} \mu \circ (f \otimes g), \qquad \mu \otimes f \otimes g \in \mathcal{E}_A^2 \otimes \mathcal{E}_A^f \otimes \mathcal{E}_A^g.$$

Proposition 3.3. *In a pre-operad C, one has*

$$\mu \circ_0 f = (-1)^f f \smile \mathrm{I}, \quad \mu \circ_1 f = -\mathrm{I} \smile f, \quad f \smile g = -(-1)^{|f|g} (\mu \circ_1 g) \circ_0 f.$$

Proof. We have

$$(-1)^f f \smile \mathrm{I} = (-1)^{f+f} (\mu \circ_0 f) \circ_f \mathrm{I} = \mu \circ_0 f, \quad -\mathrm{I} \smile f = (\mu \circ_0 \mathrm{I}) \circ_1 f = \mu \circ_1 f.$$

Also, calculate

$$\begin{aligned} f \smile g &= (-1)^f (\mu \circ_0 f) \circ_f g = (-1)^{|f||g|+f} (\mu \circ_{f-|f|} g) \circ_0 f \\ &= (-1)^{|f||g|+|f|+1} (\mu \circ_1 g) \circ_0 f = -(-1)^{|f|g} (\mu \circ_1 g) \circ_0 f, \end{aligned}$$

which is the required formula. $\square$

Lemma 3.4. *In a pre-operad C, the following composition relations hold:*

$$(f \smile g) \circ_j h = \begin{cases} (-1)^{g|h|}(f \circ_j h) \smile g & \text{if } 0 \le j \le |f|, \\ f \smile (g \circ_{j-f} h) & \text{if } f \le j \le |g| + f. \end{cases}$$

Proof. Calculate, by using the defining relations of a pre-operad:

$$\begin{aligned}
(f \smile g) \circ_j h &= (-1)^f [(\mu \circ_0 f) \circ_f g] \circ_j h \\
&= \begin{cases} (-1)^{f+|g||h|}[(\mu \circ_0 f) \circ_j h] \circ_{f+|h|} g & \text{if } 0 \le j \le |f|, \\ (-1)^f (\mu \circ_0 f) \circ_f (g \circ_{j-f} h) & \text{if } f \le j \le |g| + f, \end{cases} \\
&= \begin{cases} (-1)^{f+|g||h|}[\mu \circ_0 (f \circ_j h)] \circ_{f+|h|} g & \text{if } 0 \le j \le |f|, \\ f \smile (g \circ_{j-f} h) & \text{if } f \le j \le |g| + f, \end{cases} \\
&= \begin{cases} (-1)^{|f|+|h|+1+f+|g||h|}(f \circ_j h) \smile g & \text{if } 0 \le j \le |f|, \\ f \smile (g \circ_{j-f} h) & \text{if } f \le j \le |g| + f, \end{cases} \\
&= \begin{cases} (-1)^{g|h|}(f \circ_j h) \smile g & \text{if } 0 \le j \le |f|, \\ f \smile (g \circ_{j-f} h) & \text{if } f \le j \le |g| + f, \end{cases}
\end{aligned}$$

which is the required formula. □

4. Total composition

Definition 4.1 (cf. [4, 1]). In a pre-operad C, the *total composition* $\bullet \colon C^f \otimes C^g \to C^{f+g-1}$ is defined by

$$f \bullet g := \sum_{i=0}^{|f|} f \circ_i g \in C^{f+g-1}, \qquad |\bullet| = 0, \qquad f \otimes g \in C^f \otimes C^g.$$

The pair $\operatorname{Com} C := \{C, \bullet\}$ is called the *composition algebra* of C.

Theorem 4.2. *In a pre-operad C, one has*

$$(f \smile g) \bullet h = f \smile (g \bullet h) + (-1)^{|h|g}(f \bullet h) \smile g.$$

Proof. Use Lemma 3.4. Note that $|f \smile g| = f + g - 1$ and calculate,

$$\begin{aligned}
(f \smile g) \bullet h = \sum_{i=0}^{f+g-1} (f \smile g) \circ_i h &= \sum_{i=0}^{f-1} (f \smile g) \circ_i h + \sum_{i=f}^{f+g-1} (f \smile g) \circ_i h \\
&= (-1)^{|h|g} \sum_{i=0}^{|f|} (f \circ_i h) \smile g + \sum_{i=f}^{f+g-1} f \smile (g \circ_{i-f} h) \\
&= (-1)^{|h|g} (f \bullet h) \smile g + \sum_{i'=0}^{|g|} f \smile (g \circ_{i'} h) \\
&= (-1)^{|h|g} (f \bullet h) \smile g + f \smile (g \bullet h),
\end{aligned}$$

which is the required formula. □

Remark 4.3. This theorem tells us that *right* translations in Com C are (right) derivations of the $\smile$-algebra. It may be anticipated from Theorem 5 of [2] that the *left* translations in Com C are not derivations of the $\smile$-algebra (see the Main Theorem in Sec. 6).

5. CUP AND A PRE-COBOUNDARY OPERATOR

Definition 5.1. In a pre-operad C, define a *pre-coboundary* operator δ_μ by

$$-\delta_\mu f := [f,\mu] := f \bullet \mu - (-1)^{|f|}\mu \bullet f, \qquad \mu \otimes f \in C^2 \otimes C^f.$$

Example 5.2. In the Gerstenhaber theory [2], C is an *endomorphism* pre-operad and δ_μ is the Hochschild *coboundary operator* with the property $\delta_\mu^2 = 0$, the latter is due to the *associativity* $\mu \bullet \mu = 0$.

In this paper, we do not assume the associativity constraint.

Proposition 5.3. *In a pre-operad C, one has*

$$-\delta_\mu f = f \smile \mathrm{I} + f \bullet \mu + (-1)^{|f|}\, \mathrm{I} \smile f, \qquad \mu \otimes f \in C^2 \otimes C^f.$$

Definition 5.4. The *derivation deviation* of δ_μ over $\bullet$ is defined by

$$\operatorname{dev}_\bullet \delta_\mu (f \otimes g) := \delta_\mu (f \bullet g) - f \bullet \delta_\mu g - (-1)^{|g|}\delta_\mu f \bullet g.$$

Theorem 5.5 (cf. [2])**.** *In a pre-operad C, one has*

$$(-1)^{|g|}\operatorname{dev}_\bullet \delta_\mu (f \otimes g) = f \smile g - (-1)^{fg} g \smile f, \qquad \mu \otimes f \otimes g \in C^2 \otimes C^f \otimes C^g.$$

Proof. Two proofs can be found in [6]. ☐

6. MAIN THEOREM AND GERSTENHABER'S METHOD

In this section, we calculate the derivation deviation of the pre-coboundary operator over tribraces by using the Gerstenhaber *auxiliary variables method*. The idea of the method has been illustrated in [6].

Definition 6.1 (Gerstenhaber tribraces (cf. [2, 1]))**.** The Gerstenhaber *tribraces* $\{\cdot,\cdot,\cdot\}$ are defined as a double sum over the triangle G by

$$\{h,f,g\} := \sum_{(i,j)\in G} (h \circ_i f) \circ_j g \in C^{h+f+g-2}, \quad |\{\cdot,\cdot,\cdot\}| = 0, \quad h \otimes f \otimes g \in C^h \otimes C^f \otimes C^g.$$

Definition 6.2. The *derivation deviation* of $\delta := \delta_\mu$ over the tribraces $\{\cdot,\cdot,\cdot\}$ is defined by

$$\operatorname{dev}_{\{\cdot,\cdot,\cdot\}} \delta\,(h \otimes f \otimes g) := \delta\{h,f,g\} - \{h,f,\delta g\} - (-1)^{|g|}\{h,\delta f,g\} - (-1)^{|g|+|f|}\{\delta h,f,g\}.$$

Main Theorem (cf. [2, 5, 14])**.** *In a pre-operad C, one has*

$$(-1)^{|g|}\operatorname{dev}_{\{\cdot,\cdot,\cdot\}} \delta\,(h \otimes f \otimes g) = (h \bullet f) \smile g + (-1)^{|h|f} f \smile (h \bullet g) - h \bullet (f \smile g).$$

Notations 6.3 (auxiliary variables (cf. [2]))**.** In a pre-operad C, for $(i,j)\in G$ define

$$\begin{aligned}\Gamma_{i+1,j+1} :=& -(-1)^{|h|+|f|+|g|}\,\mathrm{I}\smile((h\circ_i f)\circ_j g)\\ &-(-1)^{|f|+|g|}\sum_{s=0}^{i-1}((h\circ_s\mu)\circ_{i+1} f)\circ_{j+1} g+(-1)^{|f|+|g|}(h\circ_i(\mathrm{I}\smile f))\circ_{j+1} g,\\ \Gamma'_{i+1,j+1} :=& +(-1)^{|g|}(h\circ_i(f\smile\mathrm{I}))\circ_{j+1} g\\ &-(-1)^{|f|+|g|}\sum_{s=i+1}^{j-f}((h\circ_s\mu)\circ_i f)\circ_{j+1} g+(-1)^{|g|}(h\circ_i f)\circ_j(\mathrm{I}\smile g),\\ \Gamma''_{i+1,j+1} :=& +(h\circ_i f)\circ_j(g\smile\mathrm{I})\\ &-(-1)^{|f|+|g|}\sum_{s=j-|f|+1}^{|h|}((h\circ_s\mu)\circ_i f)\circ_j g-((h\circ_i f)\circ_j g)\smile\mathrm{I}\,.\end{aligned}$$

Lemma 6.4. *In a pre-operad C, for $(i,j)\in G$ one has*

$$\delta((h\circ_i f)\circ_j g)-(h\circ_i f)\circ_j\delta g-(-1)^{|g|}(h\circ_i\delta f)\circ_{j+1} g=\Gamma_{i+1,j+1}+\Gamma'_{i+1,j+1}+\Gamma''_{i+1,j+1}.$$

Proof. See Appendix A. □

Notations 6.5 (truncated envelope of G')**.** Now, define a (finite) *shifted* Gerstenhaber triangle:

$$G' := \{(i,j)\in\mathbb{N}\times\mathbb{N}\,|\,1\le i\le|h|\,;\,i+f\le j\le f+|h|\}.$$

Its *envelope* is the triangle

$$G'_{env} := \{(i,j)\in\mathbb{N}\times\mathbb{N}\,|\,0\le i\le h+1\,;\,i+|f|\le j\le f+h\}=G'\sqcup\partial G'_{env}.$$

The boundary of the envelope G'_{env} is evidently $\partial G'_{env}=G'_{env}\setminus G'$. The *truncated envelope* $G'_{\widetilde{env}}$ of G' is defined by *removing* its vertices,

$$G'_{\widetilde{env}} := G'_{env}\setminus\{(|f|,0);(f+h,0);(h+1,f+h)\}=G'\sqcup\partial G'_{\widetilde{env}}.$$

Then, evidently, $G'_{\widetilde{env}}$ as having *six* vertices, turns out to be a (finite) *hexagon.*

Lemma 6.6. *In a pre-operad C, for $0\le i\le|h|$, $i+f\le j\le f+|h|$ one has*

$$(-1)^{|f|+|g|}(\delta h\circ_i f)\circ_j g=\Gamma_{ij}+\Gamma'_{i+1,j}+\Gamma''_{i+1,j+1},$$

by definition for $\Gamma_{0j},\Gamma'_{i,i+|f|}$, and $\Gamma''_{i,f+h}$ (boundary values on $\partial G'_{\widetilde{env}}$).

Proof. See Appendix B. □

Boundary Lemma. *In a pre-operad C, for $(i,j)\in\partial G'_{\widetilde{env}}$ one has*

$$\begin{aligned}\Gamma_{0j} &= (-1)^{g+|h|f} f\smile(h\circ_{j-f} g), && f\le j\le f+|h|,\\ \Gamma'_{i,i+|f|} &= (-1)^{|g|}h\circ_{i-1}(f\smile g), && 1\le i\le h,\\ \Gamma''_{i,f+h} &= (-1)^{g}(h\circ_{i-1} f)\smile g, && 1\le i\le h,\end{aligned}$$

by definition for Γ_{0f}, $\Gamma'_{h,h+|f|}$, and $\Gamma''_{1,f+h}$ (three vertex values).

Proof. See Appendix C. □

Proof of the Main Theorem (Gerstenhaber's method). First note that

$$\{h,\delta f,g\}=\sum_{i=0}^{|h|-1}\sum_{j=i+\delta f}^{|h|+|\delta f|}(h\circ_i\delta f)\circ_j g=\sum_{i=0}^{|h|-1}\sum_{j=i+f}^{|h|+|f|}(h\circ_i\delta f)\circ_{j+1} g.$$

By using Lemma 6.4 we have

$$\begin{aligned}\delta\{h,f,g\}&-\{h,f,\delta g\}-(-1)^{|g|}\{h,\delta f,g\}\\&=\sum_{i=0}^{|h|-1}\sum_{j=i+f}^{|h|+|f|}(\Gamma_{i+1,j+1}+\Gamma'_{i+1,j+1}+\Gamma''_{i+1,j+1})\\&=\sum_{i=1}^{|h|}\sum_{j=i+f}^{f+|h|}(\Gamma_{ij}+\Gamma'_{ij}+\Gamma''_{ij})=\sum_{G'}(\Gamma_{ij}+\Gamma'_{ij}+\Gamma''_{ij}).\end{aligned}$$

Now use Lemma 6.6 to see that

$$\begin{aligned}(-1)^{|f|+|g|}\{\delta h,f,g\}&=\sum_{i=0}^{|\delta h|-1}\sum_{j=i+f}^{|\delta h|+|f|}(-1)^{|f|+|g|}(\delta h\circ_i f)\circ_j g\\&=\sum_{i=0}^{|h|}\sum_{j=i+f}^{f+|h|}(\Gamma_{ij}+\Gamma'_{i+1,j}+\Gamma''_{i+1,j+1})\\&=\sum_{i=0}^{|h|}\sum_{j=i+f}^{f+|h|}\Gamma_{ij}+\sum_{i=1}^{h}\sum_{j=i+|f|}^{f+|h|}\Gamma'_{ij}+\sum_{i=1}^{h}\sum_{j=i+f}^{f+h}\Gamma''_{ij}\\&=\sum_{i=1}^{|h|}\sum_{j=i+f}^{f+|h|}\Gamma_{ij}+\sum_{j=f}^{f+|h|}\Gamma_{0j}+\sum_{i=1}^{|h|}\sum_{j=i+f}^{f+|h|}\Gamma'_{ij}+\sum_{i=1}^{h}\Gamma'_{i,i+|f|}\\&\quad+\sum_{i=1}^{|h|}\sum_{j=i+f}^{f+|h|}\Gamma''_{ij}+\sum_{i=1}^{h}\Gamma''_{i,f+h}\\&=\sum_{G'}(\Gamma_{ij}+\Gamma'_{ij}+\Gamma''_{ij})\ +\boxed{\sum_{j=f}^{f+|h|}\Gamma_{0j}+\sum_{i=1}^{h}\Gamma'_{i,i+|f|}+\sum_{i=1}^{h}\Gamma''_{i,f+h}}\,.\end{aligned}$$

One can easily see that the resulting $\boxed{\textit{boxed formula}}$ is a sum over the *boundary* $\partial G'_{\widetilde{env}}$ *of the truncated envelope* $G'_{\widetilde{env}}$ of G'. By cancelling the sums $\sum_{G'}$ and using the Boundary Lemma, we finally obtain

$$\begin{aligned}\mathrm{dev}_{\{\cdot,\cdot,\cdot\}}\,\delta\,(f\otimes g\otimes h)=&-(-1)^{g+|h|f}\sum_{j=f}^{f+|h|}f\smile(h\circ_{j-f}g)\\&-(-1)^{|g|}\sum_{i=1}^{h}h\circ_{i-1}(f\smile g)-(-1)^{g}\sum_{i=1}^{h}(h\circ_{i-1}f)\smile g\end{aligned}$$

$$= -(-1)^{g} \sum_{i=0}^{|h|} \Big[(-1)^{|h|f} f \smile (h \circ_i g) - h \circ_i (f \smile g) + (h \circ_i f) \smile g\Big]$$

$$= (-1)^{|g|} \Big[(-1)^{|h|f} f \smile (h \bullet g) - h \bullet (f \smile g) + (h \bullet f) \smile g\Big],$$

which is the required formula. □

Theorem 6.7 (cf. [2, 14])**.** *In a pre-operad C, one has*

$$(-1)^{|g|} \operatorname{dev}_{\{\cdot,\cdot,\cdot\}} \delta\, (h \otimes f \otimes g) = [h, f] \smile g + (-1)^{|h|f} f \smile [h, g] - [h, f \smile g].$$

Proof. Combine the Main Theorem with Theorem 4.2. □

Remarks 6.8. A well known first form of this theorem, found by Gerstenhaber [2] for the Hochschild *cocycles*, can be seen as a starting point of the modern *mechanical mathematics* based nowadays on the pioneering concept of (*homotopy*) *Gerstenhaber algebra* [5, 14, 13].

Our Main Theorem generalizes and modifies Gerstenhaber's Theorem 5 from [2]. One can see the appearance of the extra terms with $\delta h, \delta f$ and δg inside the tribraces. The Gerstenhaber formula is obtained when arguments in tribraces $\{h, f, g\}$ are the Hochshild *cocycles*, i. e. if $\delta h = \delta f = \delta g = 0$. As it was stated in [5, 14], on the Hochschild *cochain* level the extra terms with $\delta f, \delta g$ and δh also appear when modifying the Gerstenhaber formula. Our contribution is to work out these extra terms for an *abstract* pre-operad and give their interpretation via a *derivation deviation* of δ over the tribraces.

7. APPENDIX A

Proof of Lemma 6.4. First note that

$$\begin{aligned} -\delta_\mu((h \circ_i f) \circ_j g) =& (-1)^{|h|+|f|+|g|}\, \mathrm{I} \smile ((h \circ_i f) \circ_j g) \\ &+ \sum_{s=0}^{|h|+|f|+|g|} ((h \circ_i f) \circ_j g) \circ_s \mu + ((h \circ_i f) \circ_j g) \smile \mathrm{I}\,. \end{aligned}$$

By using the composition relations

$$((h \circ_i f) \circ_j g) \circ_s \mu = \begin{cases} (-1)^{|g|}((h \circ_i f) \circ_s \mu) \circ_{j+1} g & \text{if } 0 \le s \le j-1, \\ (h \circ_i f) \circ_j (g \circ_{s-j} \mu) & \text{if } j \le s \le j + |g|, \\ (-1)^{|g|}((h \circ_i f) \circ_{s-|g|} \mu) \circ_j g & \text{if } j + g \le s \le |h| + |f| + |g|, \end{cases}$$

cut the above sum in three pieces,

$$\begin{aligned} -\delta((h \circ_i f) \circ_j g) =& (-1)^{|h|+|f|+|g|}\, \mathrm{I} \smile ((h \circ_i f) \circ_j g) \\ &+ (-1)^{|g|} \sum_{s=0}^{j-1} ((h \circ_i f) \circ_s \mu) \circ_{j+1} g + \sum_{s=j}^{j+|g|} (h \circ_i f) \circ_j (g \circ_{s-j} \mu) \\ &+ (-1)^{|g|} \sum_{s=j+g}^{|h|+|f|+|g|} ((h \circ_i f) \circ_{s-|g|} \mu) \circ_j g + ((h \circ_i f) \circ_j g) \smile \mathrm{I}\,. \end{aligned}$$

Next, cut the first sum $\sum_{s=0}^{j-1}$ in three pieces once again by using the composition relations

$$(h \circ_i f) \circ_s \mu = \begin{cases} (-1)^{|f|}(h \circ_s \mu) \circ_{i+1} f & \text{if } 0 \leq s \leq i-1, \\ h \circ_i (f \circ_{s-i} \mu) & \text{if } i \leq s \leq i+|f|, \\ (-1)^{|f|}(h \circ_{s-|f|} \mu) \circ_i f & \text{if } i+f \leq s \leq j-1, \end{cases}$$

and desuspend summation ranges,

$$\sum_{s=j}^{j+|x|} x \circ_{s-j} \mu = \sum_{s=0}^{|x|} x \circ_s \mu = x \bullet \mu = -(-1)^{|x|}\, \mathrm{I} \smile x - \delta_\mu x - x \smile \mathrm{I}, \quad \text{for } x = f, g,$$

$$\sum_{s=i+f}^{j-1} h \circ_{s-|f|} \mu = \sum_{s=i+1}^{j-f} h \circ_s \mu,$$

$$\begin{aligned} \sum_{s=j+g}^{|h|+|f|+|g|} (h \circ_i f) \circ_{s-|g|} \mu = \sum_{s=j+1}^{|h|+|f|} (h \circ_i f) \circ_s \mu &= (-1)^{|f|} \sum_{s=j+1}^{|h|+|f|} (h \circ_{s-|f|} \mu) \circ_i f \\ &= (-1)^{|f|} \sum_{s=j-|f|+1}^{|h|} (h \circ_s \mu) \circ_i f, \end{aligned}$$

to obtain the required formula. □

8. Appendix B

8.1. **Proof of Lemma 6.6.** First note that

$$\begin{aligned} \Gamma_{ij} + \Gamma'_{i+1,j} + \Gamma''_{i+1,j+1} = {} & -(-1)^{|h|+|f|+|g|}\, \mathrm{I} \smile ((h \circ_{i-1} f) \circ_{j-1} g) \\ & - (-1)^{|f|+|g|} \sum_{s=0}^{i-2} ((h \circ_s \mu) \circ_i f) \circ_j g + (-1)^{|f|+|g|} (h \circ_{i-1} (\mathrm{I} \smile f)) \circ_j g \\ & + (-1)^{|g|} (h \circ_i (f \smile \mathrm{I})) \circ_j g - (-1)^{|f|+|g|} \sum_{s=i+1}^{j-f-1} ((h \circ_s \mu) \circ_i f) \circ_j g \\ & + (-1)^{|g|} (h \circ_i f) \circ_{j-1} (\mathrm{I} \smile g) + (h \circ_i f) \circ_j (g \smile \mathrm{I}) \\ & - (-1)^{|f|+|g|} \sum_{s=j-|f|+1}^{|h|} ((h \circ_s \mu) \circ_i f) \circ_j g - ((h \circ_i f) \circ_j g) \smile \mathrm{I}\,. \end{aligned}$$

We must compare it term by term with

$$\begin{aligned} -(\delta h \circ_i f) \circ_j g = {} & \Bigg(\bigg((-1)^{|h|}\, \mathrm{I} \smile h + \sum_{s=0}^{|h|} h \circ_s \mu + h \smile \mathrm{I} \bigg) \circ_i f \Bigg) \circ_j g \\ = {} & (-1)^{|h|} ((\mathrm{I} \smile h) \circ_i f) \circ_j g + \sum_{s=0}^{i-2} ((h \circ_s \mu) \circ_i f) \circ_j g + ((h \circ_{i-1} \mu) \circ_i f) \circ_j g \\ & + ((h \circ_i \mu) \circ_i f) \circ_j g + \sum_{s=i+1}^{j-f-1} ((h \circ_s \mu) \circ_i f) \circ_j g + ((h \circ_{j-f} \mu) \circ_i f) \circ_j g \end{aligned}$$

$$+((h\circ_{j-|f|}\mu)\circ_i f)\circ_j g+\sum_{s=j-|f|+1}^{|h|}((h\circ_s\mu)\circ_i f)\circ_j g+((h\smile \mathrm{I})\circ_i f)\circ_j g.$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note the ***ground identities***

$$\begin{aligned}
((\mathrm{I}\smile h)\circ_i f)\circ_j g &= ((\mathrm{I}\smile (h\circ_{i-1} f))\circ_j g=\mathrm{I}\smile ((h\circ_{i-1} f)\circ_{j-1} g),\\
(h\circ_{i-1}\mu)\circ_i f &= h\circ_{i-1}(\mu\circ_1 f)=-h\circ_{i-1}(\mathrm{I}\smile f),\\
(h\circ_i\mu)\circ_i f &= h\circ_i(\mu\circ_0 f)=(-1)^f h\circ_i(f\smile \mathrm{I}),\\
((h\circ_{j-f}\mu)\circ_i f)\circ_j g &=(-1)^{|f|}((h\circ_i f)\circ_{j-1}\mu)\circ_j g=(-1)^{|f|}(h\circ_i f)\circ_{j-1}(\mu\circ_1 g)\\
&=-(-1)^{|f|}(h\circ_i f)\circ_{j-1}(\mathrm{I}\smile g),\\
((h\circ_{j-|f|}\mu)\circ_i f)\circ_j g &=(-1)^{|f|}((h\circ_i f)\circ_j\mu)\circ_j g=(-1)^{|f|}(h\circ_i f)\circ_j(\mu\circ_0 g)\\
&=-(-1)^{|f|+|g|}(h\circ_i f)\circ_j(g\smile \mathrm{I}),\\
((h\smile \mathrm{I})\circ_i f)\circ_j g &= (-1)^{|f|}((h\circ_i f)\smile \mathrm{I})\circ_j g=(-1)^{|f|+|g|}((h\circ_i f)\circ_j g)\smile \mathrm{I},
\end{aligned}$$

which prove the required formula. $\square$

8.2. **Proposition/recapitulation.** *In a pre-operad C, for $(i,j)\in G'$ the auxiliary variables read*

$$\Gamma_{ij}=-(-1)^{|h|+|f|+|g|}((\mathrm{I}\smile h)\circ_i f)\circ_j g-(-1)^{|f|+|g|}\sum_{s=0}^{i-1}((h\circ_s\mu)\circ_i f)\circ_j g,$$

$$\Gamma'_{ij}=-(-1)^{|f|+|g|}\sum_{s=i-1}^{j-f}((h\circ_s\mu)\circ_{i-1} f)\circ_j g,$$

$$\Gamma''_{ij}=-(-1)^{|f|+|g|}\sum_{s=j-f}^{|h|}((h\circ_s\mu)\circ_{i-1} f)\circ_{j-1} g-(-1)^{|f|+|g|}((h\smile \mathrm{I})\circ_{i-1} f)\circ_{j-1} g.$$

Proof. Use the ***ground identities*** from the previous subsection 8.1. $\square$

9. Appendix C

Proof of the Boundary Lemma. First prove the

Vertex Proposition. In a pre-operad C, one has

$$\begin{aligned}
\Gamma_{0,f+|h|} &= (-1)^{g+|h|f} f\smile (h\circ_{|h|} g),\\
\Gamma'_{1f} &= (-1)^{|g|} h\circ_0 (f\smile g),\\
\Gamma''_{h,f+h} &= (-1)^{g}(h\circ_{|h|} f)\smile g.
\end{aligned}$$

Proof. We calculate these vertex values in a standard way, by using Lemma 6.6. First calculate $\Gamma_{0,f+|h|}$. Use Lemma 6.6 and $\Gamma''_{1,f+h}$ from the Boundary Lemma to note that

$$\Gamma_{0,f+|h|}+\Gamma'_{1,f+|h|}+\Gamma''_{1,f+h}=\Gamma_{0,f+|h|}-(-1)^{|f|+|g|}\sum_{s=0}^{|h|}((h\circ_s\mu)\circ_0 f)\circ_{f+|h|} g$$

$$+ (-1)^{g}(h \circ_0 f) \smile g.$$

We must compare it term by term with

$$\begin{aligned} -(\delta h \circ_0 f) \circ_{f+|h|} g &= \left(\left((-1)^{|h|}\, \mathrm{I} \smile h + \sum_{s=0}^{|h|} h \circ_s \mu + h \smile \mathrm{I}\right) \circ_0 f\right) \circ_{f+|h|} g \\ &= (-1)^{|h|}((\mathrm{I} \smile h) \circ_0 f) \circ_{f+|h|} g + \sum_{s=0}^{|h|}((h \circ_s \mu) \circ_0 f) \circ_{f+|h|} g \\ &\quad + ((h \smile \mathrm{I}) \circ_0 f) \circ_{f+|h|} g. \end{aligned}$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note that

$$\begin{aligned} &(-1)^{|h|}((\mathrm{I} \smile h) \circ_0 f) \circ_{f+|h|} g = (-1)^{|h|+|f|h}((\mathrm{I} \circ_0 f) \smile h) \circ_{f+|h|} g \\ &= (-1)^{|h|f+|f|}(f \smile h) \circ_{f+|h|} g = -(-1)^{|g|+|f|} \boxed{(-1)^{g+|h|f} f \smile (h \circ_{|h|} g)}, \\ &((h \smile \mathrm{I}) \circ_0 f) \circ_{f+|h|} g = (-1)^{|f|}((h \circ_0 f) \smile \mathrm{I}) \circ_{f+|h|} g = (-1)^{|f|}(h \circ_0 f) \smile (\mathrm{I} \circ_0 g) \\ &= (-1)^{|f|}(h \circ_0 f) \smile g, \end{aligned}$$

which lead one to the required formula for $\Gamma_{0,f+|h|}$.

Next calculate Γ'_{1f}. Use Lemma 6.6 and $\Gamma_{0,f}$ from the Boundary Lemma to note that

$$\begin{aligned} \Gamma_{0f} + \Gamma'_{1f} + \Gamma''_{1,f+1} &= (-1)^{f|h|+g} f \smile (h \circ_0 g) + \Gamma'_{1f} \\ &\quad - (-1)^{|f|+|g|} \sum_{s=1}^{|h|}((h \circ_s \mu) \circ_0 f) \circ_f g - (-1)^{|f|+|g|}((h \smile \mathrm{I}) \circ_0 f) \circ_f g. \end{aligned}$$

We must compare it term by term with

$$\begin{aligned} -(\delta h \circ_0 f) \circ_f g &= \left(\left((-1)^{|h|}\, \mathrm{I} \smile h + \sum_{s=0}^{|h|} h \circ_s \mu + h \smile \mathrm{I}\right) \circ_0 f\right) \circ_f g \\ &= (-1)^{|h|}((\mathrm{I} \smile h) \circ_0 f) \circ_f g + ((h \circ_0 \mu) \circ_0 f) \circ_f g \\ &\quad + \sum_{s=1}^{|h|}((h \circ_s \mu) \circ_0 f) \circ_f g + ((h \smile \mathrm{I}) \circ_0 f) \circ_f g. \end{aligned}$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note that

$$\begin{aligned} ((\mathrm{I} \smile h) \circ_0 f) \circ_f g &= (-1)^{|f|h}((\mathrm{I} \circ_0 f) \smile h) \circ_f g = (-1)^{|f|h}(f \smile h) \circ_f g \\ &= (-1)^{|f|h} f \smile (h \circ_0 g), \\ ((h \circ_0 \mu) \circ_0 f) \circ_f g &= (h \circ_0 (\mu \circ_0 f)) \circ_f g = h \circ_0 ((\mu \circ_0 f) \circ_f g) \\ &= -(-1)^{|f|+|g|} \boxed{(-1)^{|g|} h \circ_0 (f \smile g)}, \end{aligned}$$

which lead one to the required formula for Γ'_{1f}.

At last calculate $\Gamma''_{h,f+h}$. Use Lemma 6.6 and $\Gamma'_{h,f+|h|}$ from the Boundary Lemma to note that

$$\Gamma_{|h|,f+|h|} + \Gamma'_{h,f+|h|} + \Gamma''_{h,f+h} = -(-1)^{|f|+|g|+|h|}((\mathrm{I} \smile h) \circ_{|h|} f) \circ_{f+|h|} g$$

$$- (-1)^{|f|+|g|}\sum_{s=0}^{|h|-1}((h\circ_s\mu)\circ_{|h|} f)\circ_{f+|h|} g + (-1)^{|g|}h\circ_{|h|}(f\smile g) + \Gamma''_{h,f+h}.$$

We must compare it term by term with

$$\begin{aligned} -(\delta h\circ_{|h|} f)\circ_{f+|h|} g &= \Bigl(\Bigl((-1)^{|h|}\,\mathrm{I}\smile h + \sum_{s=0}^{|h|} h\circ_s\mu + h\smile \mathrm{I}\Bigr)\circ_{|h|} f\Bigr)\circ_{f+|h|} g \\ &= (-1)^{|h|}((\mathrm{I}\smile h)\circ_{|h|} f)\circ_{f+|h|} g + \sum_{s=0}^{|h|-1}((h\circ_s\mu)\circ_{|h|} f)\circ_{f+|h|} g \\ &\quad + ((h\circ_{|h|}\mu)\circ_{|h|} f)\circ_{f+|h|} g + ((h\smile \mathrm{I})\circ_{|h|} f)\circ_{f+|h|} g. \end{aligned}$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note that

$$\begin{aligned} ((h\circ_{|h|}\mu)\circ_{|h|} f)\circ_{f+|h|} g &= (h\circ_{|h|}(\mu\circ_0 f))\circ_{f+|h|} g = h\circ_{|h|}((\mu\circ_0 f)\circ_f g) \\ &= (-1)^f h\circ_{|h|}(f\smile g), \end{aligned}$$

$$\begin{aligned} ((h\smile \mathrm{I})\circ_{|h|} f)\circ_{f+|h|} g &= (-1)^{|f|}((h\circ_{|h|} f)\smile \mathrm{I})\circ_{f+|h|} g \\ &= (-1)^{|f|}(h\circ_{|h|} f)\smile(\mathrm{I}\circ_0 g) = -(-1)^{|f|+|g|}\boxed{(-1)^g(h\circ_{|h|} f)\smile g}, \end{aligned}$$

which lead one to the required formula for $\Gamma''_{h,f+h}$. □

Now, when the vertex values found, we can accomplish the proof of the Boundary Lemma. First calculate Γ_{0j}. Use Lemma 6.6 to note that

$$\begin{aligned} \Gamma_{0j} + \Gamma'_{1j} + \Gamma''_{1,j+1} &= \Gamma_{0j} - (-1)^{|f|+|g|}\sum_{s=0}^{j-f}((h\circ_s\mu)\circ_0 f)\circ_j g \\ &\quad - (-1)^{|f|+|g|}\sum_{s=j-|f|}^{|h|}((h\circ_s\mu)\circ_0 f)\circ_j g - (-1)^{|f|+|g|}((h\smile \mathrm{I})\circ_0 f)\circ_j g. \end{aligned}$$

We must compare it term by term with

$$\begin{aligned} -(\delta h\circ_0 f)\circ_j g &= \Bigl(\Bigl((-1)^{|h|}\,\mathrm{I}\smile h + \sum_{s=0}^{|h|} h\circ_s\mu + h\smile \mathrm{I}\Bigr)\circ_0 f\Bigr)\circ_j g \\ &= (-1)^{|h|}((\mathrm{I}\smile h)\circ_0 f)\circ_j g + \sum_{s=0}^{j-f}((h\circ_s\mu)\circ_0 f)\circ_j g \\ &\quad + \sum_{s=j-|f|}^{|h|}((h\circ_s\mu)\circ_0 f)\circ_j g + ((h\smile \mathrm{I})\circ_0 f)\circ_j g. \end{aligned}$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note that

$$\begin{aligned} (-1)^{|h|}((\mathrm{I}\smile h)\circ_0 f)\circ_j g &= (-1)^{|h|+h|f|}((\mathrm{I}\circ_0 f)\smile h)\circ_j g \\ = (-1)^{|h|f+|f|}(f\smile h)\circ_j g &= -(-1)^{|f|+|g|}\boxed{(-1)^{g+|h|f} f\smile(h\circ_{j-f} g)}, \end{aligned}$$

which leads one to the required formula for Γ_{0j}.

Next calculate $\Gamma'_{i,i+|f|}$. Use Lemma 6.6 to note that

$$\begin{aligned}\Gamma_{i-1,i+|f|}+\Gamma'_{i,i+|f|}+\Gamma''_{i,i+f}&=-(-1)^{|f|+|g|+|h|}((\mathrm{I}\smile h)\circ_{i-1}f)\circ_{i+|f|}g\\&-(-1)^{|f|+|g|}\sum_{s=0}^{i-2}((h\circ_s\mu)\circ_{i-1}f)\circ_{i+|f|}g+\Gamma'_{i,i+|f|}\\&-(-1)^{|f|+|g|}\sum_{s=i}^{|h|}((h\circ_s\mu)\circ_{i-1}f)\circ_{i+|f|}g\\&-(-1)^{|f|+|g|}((h\smile\mathrm{I})\circ_{i-1}f)\circ_{i+|f|}g.\end{aligned}$$

We must compare it term by term with

$$\begin{aligned}-(\delta h\circ_{i-1}f)\circ_{i+|f|}g&=\left(\left((-1)^{|h|}\,\mathrm{I}\smile h+\sum_{s=0}^{|h|}h\circ_s\mu+h\smile\mathrm{I}\right)\circ_{i-1}f\right)\circ_{i+|f|}g\\&=(-1)^{|h|}((\mathrm{I}\smile h)\circ_{i-1}f)\circ_{i+|f|}g+\sum_{s=0}^{i-2}((h\circ_s\mu)\circ_{i-1}f)\circ_{i+|f|}g\\&\quad+((h\circ_{i-1}\mu)\circ_{i-1}f)\circ_{i+|f|}g+\sum_{s=i}^{|h|}((h\circ_s\mu)\circ_{i-1}f)\circ_{i+|f|}g\\&\quad+((h\smile\mathrm{I})\circ_{i-1}f)\circ_{i+|f|}g.\end{aligned}$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note that

$$\begin{aligned}((h\circ_{i-1}\mu)\circ_{i-1}f)\circ_{i+|f|}g&=(h\circ_{i-1}(\mu\circ_0 f))\circ_{i+|f|}g=h\circ_{i-1}((\mu\circ_0 f)\circ_f g)\\&=-(-1)^{|f|+|g|}\boxed{(-1)^{|g|}h\circ_{i-1}(f\smile g)},\end{aligned}$$

which leads one to the required formula for $\Gamma'_{i,i+|f|}$.

At last calculate $\Gamma''_{i,f+h}$. Use Lemma 6.6 to note that

$$\begin{aligned}\Gamma_{i-1,f+|h|}+\Gamma'_{i,f+|h|}+\Gamma''_{i,f+h}&=-(-1)^{|f|+|g|+|h|}((\mathrm{I}\smile h)\circ_{i-1}f)\circ_{f+|h|}g\\&-(-1)^{|f|+|g|}\sum_{s=0}^{i-2}((h\circ_s\mu)\circ_{i-1}f)\circ_{f+|h|}g\\&-(-1)^{|f|+|g|}\sum_{s=i-1}^{|h|}((h\circ_s\mu)\circ_{i-1}f)\circ_{f+|h|}g+\Gamma''_{i,f+h}.\end{aligned}$$

We must compare it term by term with

$$\begin{aligned}-(\delta h\circ_{i-1}f)\circ_{f+|h|}g&=\left(\left((-1)^{|h|}\,\mathrm{I}\smile h+\sum_{s=0}^{|h|}h\circ_s\mu+h\smile\mathrm{I}\right)\circ_{i-1}f\right)\circ_{f+|h|}g\\&=(-1)^{|h|}((\mathrm{I}\smile h)\circ_{i-1}f)\circ_{f+|h|}g+\sum_{s=0}^{i-2}((h\circ_s\mu)\circ_{i-1}f)\circ_{f+|h|}g\end{aligned}$$

$$+\sum_{s=i-1}^{|h|}((h\circ_s\mu)\circ_{i-1}f)\circ_{f+|h|}g+((h\smile \mathrm{I})\circ_{i-1}f)\circ_{f+|h|}g.$$

Now, recall the sign $(-1)^{|f|+|g|}$ and use composition relations to note that

$$\begin{aligned}((h\smile \mathrm{I})\circ_{i-1}f)\circ_{f+|h|}g&=(-1)^{|f|}((h\circ_{i-1}f)\smile \mathrm{I})\circ_{f+|h|}g\\&=(-1)^{|f|}(h\circ_{i-1}f)\smile(\mathrm{I}\circ_0 g)=-(-1)^{|f|+|g|}\boxed{(-1)^{g}(h\circ_{i-1}f)\smile g},\end{aligned}$$

which leads one to the required formula for $\Gamma''_{i,f+h}$. ☐

Acknowledgements

One of the authors (E. Paal) would like to thank Jim Stasheff for reading of the preliminary manuscript and useful discussions and suggestions.

References

- [1] V. Coll, M. Gerstenhaber, and S. D. Schack Universal deformation formulas and breaking symmetry. J. Pure Appl. Algebra, 1993, **90**, 201-219.
- [2] M. Gerstenhaber. The cohomology structure of an associative ring. Ann. of Math. 1963, **78**, 267-288.
- [3] M. Gerstenhaber. On the deformation theory of rings and algebras: III. Ann. of Math. 1968, **88**, 1-34.
- [4] M. Gerstenhaber, A. Giaquinto, and S. D. Schack Algebras, bialgebras, quantum groups, and algebraic deformations. In *Deformation Theory and Quantum Groups with Applications to Mathematical Physics, Amherst, MA, 1990.* M. Gerstenhaber and J. Stasheff, Eds. Contemp. Math. 1992, **134**, 51-92.
- [5] M. Gerstenhaber and A. A. Voronov. Homotopy G-algebras and moduli space operad. Int. Math. Res. Notices, 1995, **3**, 141-153; hep-th/9409063.
- [6] L. Kluge, E. Paal, and J. Stasheff. Invitation to composition. Comm. Algebra, 2000, **28**, 1405-1422; math. QA/9908083.
- [7] J. L. Loday, J. Stasheff, and A. A. Voronov, Eds. *Operads: Proceedings of Renaissance Conferences.* Contemp. Math. 1997, **202**.
- [8] M. Markl. Simplex, associahedron, and cyclohedron. Contemp. Math. 1999, **227**, 235-265; alg-geom/9707009.
- [9] J. P. May. *The Geometry of Iterated Loop Spaces.* Lecture Notes in Math. 1972, **271**.
- [10] S. Shnider and S. Sternberg *Quantum Groups from Coalgebras to Drinfeld Algebras.* Int. Press Inc.: Boston, 1994.
- [11] J. D. Stasheff. Homotopy associativity of H-spaces, I, II. Trans. Amer. Math. Soc. 1963, **108**, 275-292, 293-312.
- [12] J. Stasheff. From operads to ‘physically’ inspired theories. Contemp. Math. 1997, **202**, 53-81.
- [13] A. A. Voronov. Homotopy Gerstenhaber algebras; math. QA/9908040.
- [14] A. A. Voronov and M. Gerstenhaber. Higher-order operations on the Hochschild complex. Funktsinal. Anal. i Prilozhen. 1995, **29**, 1-6 (in Russian).

Department of Mathematics, Tallinn Technical University
Ehitajate tee 5, 19086 Tallinn, Estonia
e-mails: liivi.kluge.ttu@mail.ee and epaal@edu.ttu.ee